



# Newsletter

[www.carovail.com](http://www.carovail.com)  
[www.facebook.com/CaroVail](http://www.facebook.com/CaroVail)

CaroVail

Agronomy Update

August 1, 2016

## Locations

### *Auburn*

55 Columbus St  
Auburn, NY 13021  
315-253-7379

### *Bernardston*

472 Northfield Road  
Bernardston, MA  
01337  
413-648-9900

### *Niverville*

831 Route 28  
Niverville, NY 12130  
518-784-9166

### *Oriskany Falls*

8341 US State Rt 20  
Oriskany Falls, NY  
13425  
315-841-3201

### *Salem*

4134 State Rt 22  
Salem, NY 12865  
518-854-9446

### *Tri Valley Crop Ctr*

337 State Hwy 162  
Sprakers, NY 12166  
518-673-5336

## **SPECIAL EDITION**

### **Drought Conditions**

Frank Flis

Many areas in the northeast are experiencing moisture challenges to varying degrees. Having adequate forage supplies for the coming year may be a major concern for growers in some areas. Care in managing fields to ensure best possible results may take planning and unemotional action. Calm, rational analysis will be required.

Maximizing potential while preserving crop integrity is the challenge, especially with perennial crops like alfalfas, grasses, and pastures. Getting a handle on inventories both on hand and needed for the coming year is an important part of the process. Do you know how much you have and how much you need?

Please take time to consult your advisors on recommended next steps. Speak with your nutritionist on needed forage extenders and your crop advisors on field management. In this special addition of CaroVail's newsletter is some advice and experience from The University of Wisconsin. As always, CaroVail personnel are available to assist.

Good management of moisture stressed forages will increase maintenance of a good stand and reduce the loss of forage production. Management of alfalfa, grass hayfields, pasture and corn each have some special considerations.

## Alfalfa

Moisture stress has the following effects on the alfalfa plant:

- Cell enlargement is inhibited.
- The number of basal buds and the number of shoots or stems/plant is reduced when moisture stress occurs in the first 14 days after a harvest.
- The stem internode length is reduced; thus the flowering is seen at reduced plant height.
- Leaf area/leaf size and leaf growth rate is reduced, although to a lesser degree than stem growth. Therefore leaf to stem ratio is higher under moisture stress.
- Stem nitrogen percentage is increased while leaf nitrogen percentage is decreased, therefore whole plant nitrogen (CP) may be reduced though effect varies with severity/timing of moisture stress.
- NDF is generally decreased, though effect varies with severity/timing of moisture stress.

### Recommendations:

1. If stand is over 10 inches tall and flowering, harvest if economic to do so. Moisture stressed alfalfa should be mowed at the normal cutting height. There is no advantage to raising the cutting height. Alfalfa can regrow from axillary buds higher up on the stubble but they are smaller and produce lower yield than stems growing from the crown buds. Since quality is not declining as rapidly with advancing maturity as under normal growing conditions, let the plants approach 100% bloom before harvest to allow the plant to build nonstructural carbohydrate reserves.
2. If stand is 10 inches or less tall and flowering, do not cut. Let regrowth come through existing growth. Mowing will not increase regrowth.
3. Make sure that soil fertility is at optimum levels.
4. Scout and control potato leaf hopper, army worm and other insects.
5. New seedings should not be harvested during the season but may be harvested in late August if adequate growth is present to harvest. A late fall cutting may also be taken. The key is to time harvests so that alfalfa either has no regrowth or more than 8 inches of regrowth at frost.

## Grassy hay fields

Moisture stress effects on grasses are similar to alfalfa except that, if stems are present, forage quality of grasses may be lowered by stress rather than increased. Most grassy fields are stunted but are leafy and have few stems.

### Recommendations:

1. Harvest if tonnage justifies and/or height is over 8 to 10 inches.
2. Apply 40 lb/a nitrogen to stimulate fall growth if rain occurs before mid-August. This cannot be manure since it will become available too slowly to provide optimum fall growth.

## Pasture

Most pastures are short but regrowth will occur when it rains if adequate nitrogen is present.

### Recommendations:

1. Graze to take advantage of any existing vegetation.
2. Mow tall, weedy or brushy growth.
3. Apply 40 lb/a nitrogen to stimulate growth as soon after Aug 1 as possible.

## Corn for forage

Many fields are stunted, some have significant firing.

### Recommendations:

1. Plants will put out more growth, all are too wet to ensile now. Check moisture before chopping or bailing to ensure excessive moisture does not cause poor fermentation and seepage losses.
2. If grazing, consider potential for nitrate toxicity. This is especially likely to be a problem if growth was reduced to less than 50% of normal and/or high levels of nitrogen were applied. Samples taken for nitrate test must be frozen or analyzed immediately as nitrate will decline in tissue over 3 to 4 hours. If above toxic, levels feed hay or some other forage in the morning and graze corn a couple hours in the afternoon.

**Auburn**

**Bernardston**

**Niverville**

**Oriskany Falls**

**Salem**

**Tri Valley**

**Auburn**

**Bernardston**

**Niverville**

**Oriskany Falls**

**Salem**

**Tri Valley**

