

---

## SUPPLEMENTAL LABELING

---

### RESTRICTED USE PESTICIDE DUE TO ACUTE TOXICITY

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

#### Syngenta Crop Protection, LLC

P. O. Box 18300

Greensboro, North Carolina 27419-8300

SCP 1431A-S16 1211

#### Gramoxone® SL 2.0

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide

Vegetables (Cucumber, Melons, Pumpkin, Squash) – For Use Only in the State of New Jersey

Active Ingredient:

Paraquat dichloride (1,1'-dimethyl-4,4'-bipyridinium dichloride)..... 30.1%

Other Ingredients: ..... 69.9%

Total: ..... 100.0%

Contains 2.0 pounds paraquat cation per gallon as 2.762 pounds salt per gallon.

Contains alerting agent (odor), emetic, dye.

EPA Reg. No. 100-1431

**KEEP OUT OF REACH OF CHILDREN.**



**DANGER / POISON**  
**PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

- **NEVER PUT INTO FOOD, DRINK OR OTHER CONTAINERS.**
- **IF SWALLOWED, TAKE IMMEDIATE ACTION AS PRESCRIBED IN FIRST AID STATEMENT. SYMPTOMS ARE PROLONGED AND PAINFUL.**
- **DO NOT USE OR STORE IN OR AROUND THE HOME.**
- **DO NOT REMOVE CONTENTS EXCEPT FOR IMMEDIATE USE.**
- **THE ODOR OF THIS PRODUCT IS FROM THE ALERTING AGENT WHICH HAS BEEN ADDED, NOT FROM PARAQUAT.**

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using GRAMOXONE SL 2.0 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**syngenta**

## DIRECTIONS FOR USE

Crop	Maximum Number of Applications	Use Pattern	Gramoxone SL 2.0 Rate/Acre	Minimum Total Spray per Acre	Grazing or Preharvest Interval	Precautions, Restrictions and Comments
<b>VEGETABLES</b>  Cucumber Melons Pumpkin Squash  For Use Only in the State of New Jersey	3	Directed/ Shielded Spray	1.95 pts.	Ground: 20 gals.	14 days	<ul style="list-style-type: none"> <li>For control or suppression of emerged weeds between rows after crop establishment.</li> <li>Use precision directed spray equipment adjusted to prevent spray contact with crop plants.</li> <li>Add either: 1) A non-ionic surfactant containing 75% or more surface active agent at 0.25% v/v (2 pts. per 100 gals.), of the finished spray volume for ground applications, <b>OR</b> 2) Crop Oil Concentrate (non-phytotoxic) containing 15-20% approved emulsifier, at 1.0% v/v (1 gal. per 100 gals.) of the finished spray volume.</li> <li>Apply when weeds are 6" or less in height and actively growing.</li> <li>Do not exceed 30 psi nozzle pressure or spray under conditions which may cause excessive drift.</li> <li>Do not allow spray to contact green stems or foliage, as injury may result.</li> <li>Do not use on soils with less than 5% clay.</li> <li>Do not allow animals to graze in treated areas.</li> <li>Do not harvest within 14 days after application.</li> <li>Do not apply after bloom.</li> <li>Do not apply this product through any type of irrigation system.</li> </ul>

Gramoxone® SL 2.0 and the Syngenta logo trademark of a Syngenta Group Company

©2012 Syngenta

GRAMOXONE SL 2.0 1431A-S16 1211 – tb – 02/08/12