

Product Bulletin



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Indar[®] 2F

EPA Reg. No. 62719-416

2(ee) Recommendation [†]

Control of Exobasidium leaf and fruit spot in Blueberry

(For Distribution and Use Only in the States of AL, AR, CT, DE, FL, GA, IA, IL, IN, KY, LA, MA, MD, ME, MI, MN, MO, MS, NC, NH, NJ, OH, PA, RI, SC, TN, VA, VT, WI, and WV)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Indar[®] 2F fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Indar 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Indar 2F.

Directions for Use

Refer to product label for Mixing, Handling and Application instructions.

Indar 2F is primarily a protectant fungicide and the best disease management results against Exobasidium leaf and fruit spot of blueberry are achieved when Indar 2F is applied prior to disease infection.

Use 6 fl oz of Indar 2F (0.094 lb active) per acre in a minimum of 20 gallons of water by ground or 10 gallons of water by air.

Specific Use Restrictions:

- Do not make more than 4 applications or apply more than 24 fl oz of Indar 2F (0.75 lb active) per acre per year.
- **Preharvest Interval:** Do not apply within 30 days of harvest.
- Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.

[®]Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R349-017

Issued: 03/18/14

Initial printing.