

Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Indar[®] 2F

EPA Reg. No. 62719-416

2(ee) Recommendation[†]

(For Use and Distribution in the States of Massachusetts, New Jersey, and Rhode Island)

Suppression of Fairyring Complex in Cranberry

ATTENTION

[†] This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Indar[®] 2F fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Except as specified in this product bulletin, use of Indar 2F is subject to all use precautions and limitations imposed by the label affixed to the container for Indar 2F.

Refer to product label for Resistance Management, Mixing and Application instructions.

Disease	Indar 2F (fl oz/acre)	Use Directions	Restrictions
fairyring complex (suppression)	12 (0.188 lb active)	Measure ring diameter and add 10 ft to the diameter. Begin applications at bud break and repeat once if necessary 14 days later. Apply in 30 to 100 gallons of water to the affected area. Irrigation for 1 to 2 hours following application is advisable to ensure penetration to the base of the plant.	<ul style="list-style-type: none">• Do not make more than 4 applications or apply more than 48 fl oz of Indar 2F (0.75 lb active) per acre per year.• Preharvest Interval: Do not apply within 30 days of harvest.

[®]Trademark of Dow AgroSciences LLC

R349-005

Issued: 02/19/08

Initial printing.