

Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Radiant[®] SC

EPA Reg. No. 62719-545

Special 2(ee) Recommendation[†]

(For Distribution and Use Only Within New York State)

Suppression of Spotted Wing Drosophila on Strawberry

ATTENTION

[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Radiant[®] SC insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Radiant SC according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Radiant SC.

Directions for Use

Refer to product label for Radiant SC for General Use Precautions, Mixing Directions and Application Directions.

Radiant SC can be used as part of an integrated program to manage spotted wing drosophila. This use is limited to ground application.

Apply 6 to 10 fl oz of Radiant SC per acre to strawberry for suppression of spotted wing drosophila. Use a higher rate in the rate range for moderate to severe infestations and/or larger plant volume. Begin applications at first signs of adult activity. Due to the occurrence of multiple generations in a growing season, repeated applications may be required. Follow resistance management recommendations on the product label.

Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area. Refer to specific crop sections of the product label for additional information and restrictions.

[®]Trademark of Dow AgroSciences LLC

R358-017

Issued: 11/28/11

Initial printing.