

Supplemental Label



EPA Reg. No. 59639-132
(Except California and New York)

APPLICATION OF SELECT MAX[®] HERBICIDE WITH INSIDE TECHNOLOGY[™] IN OKRA AND TREE NUTS CROP GROUP 14-12

This supplemental label expires May 23, 2021 and must not be used or distributed after this date.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR *SELECT MAX HERBICIDE WITH INSIDE TECHNOLOGY* BEFORE APPLYING. USE OF *SELECT MAX HERBICIDE WITH INSIDE TECHNOLOGY* ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR *SELECT MAX HERBICIDE WITH INSIDE TECHNOLOGY*.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Restriction: Do not plant rotational crops until 30 days after application of *Select MAX Herbicide with Inside Technology* unless the crop is listed on *Select MAX Herbicide with Inside Technology* label.

CROP SPECIFIC USE DIRECTIONS AND RESTRICTIONS FOR *SELECT MAX HERBICIDE WITH INSIDE TECHNOLOGY* HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
Okra	3 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) at 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions <ul style="list-style-type: none">• Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application.• Do not make more than 4 applications per acre per year.• Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.

continued

CROP SPECIFIC USE DIRECTIONS AND RESTRICTIONS FOR SELECT MAX HERBICIDE WITH INSIDE TECHNOLOGY HERBICIDE

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Annual Grass Use Rate Per Acre ⁽²⁾	Perennial Grass Use Rate Per Acre ⁽²⁾	Adjuvant Recommendation ⁽³⁾	Ammonium Sulfate Recommendation ⁽⁴⁾	Special Use Instructions And Restrictions
Tree Nuts Crop Group 14-12 including: African Nut-tree Almond Beechnut Brazil Nut Brazilian Pine Bunya Bur Oak Butternut Cajou Nut Candlenut Cashew Chestnut Chinquapin Coconut Coquito Nut Dika Nut Ginkgo Guiana Chestnut Hazelnut (Filbert) Heartnut Hickory Nut Japanese Horse-chestnut Macadamia Nut Mongongo Nut Monkey-pot Monkey Puzzle Nut Okari Nut Pachira Nut Peach Palm Nut Pecan Pequi Pili Nut Pine Nut Pistachio Sapucaia Nut Tropical Almond Walnut, Black Walnut, English Yellowhorn Cultivars, varieties, and/or hybrids of these.	14 days	9 to 16 fl oz	12 to 16 fl oz	Non-ionic Surfactant (NIS) At 0.25% v/v	None	For repeat applications make on a minimum of a 14 day interval. Restrictions <ul style="list-style-type: none"> Do not apply more than 16 fl oz/A (0.121 lb ai/A) per application. Do not make more than 4 applications per acre per year. Do not apply more than 64 fl oz/A (0.485 lb ai/A) per year.

⁽¹⁾ Select MAX Herbicide with Inside Technology is not for use on vegetable crops being grown for seed production unless specific use directions are provided.

⁽²⁾ See annual and perennial grass control tables for specific use rates.

⁽³⁾ Non-ionic surfactant (NIS) in this case refers to an adjuvant containing at least 80% non-ionic surfactant. Crop oil concentrate in this case refers to both crop oil concentrate and crop oil concentrate blends. **Acceptable crop oil concentrates** would be those that contain a minimum of 80% oils and 15% emulsifier. **Acceptable crop oil concentrate blends** would be those that contain a minimum of 60% oils and 25-40% surfactants and emulsifiers. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils.

⁽⁴⁾ Use spray grade ammonium sulfate. The use of ammonium sulfate does not take the place of the required adjuvant.

USE DIRECTIONS FOR ANNUAL GRASSES ALL CROPS

- Apply only to actively growing grasses at specified weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the specified growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

Restrictions

- Do not exceed the maximum per application rate listed in CROP SPECIFIC USE DIRECTIONS AND RESTRICTIONS FOR *Select MAX Herbicide with Inside Technology*.
- Do not exceed the maximum yearly rate listed in Table 1, CROP SPECIFIC USE DIRECTIONS AND RESTRICTIONS FOR *Select MAX Herbicide with Inside Technology*.

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT* (inches)	APPLICATION RATES	
			MINIMUM RATE fl oz/A	MAXIMUM RATE ⁽¹⁾ fl oz/A
Barnyardgrass	<i>Echinochloa crus-galli</i>	2 to 8	9	16
Broadleaf Signalgrass	<i>Brachiaria platyphylla</i>	2 to 6	9	16
Brome				
California	<i>Bromus carinatus</i>	2 to 6	9	16
Cheat	<i>Bromus secalinus</i>	2 to 6	9	16
Downy	<i>Bromus tectorum</i>	2 to 6	9	16
Ripgut	<i>Bromus diandrus</i>	2 to 6	9	16
Canarygrass	<i>Phalaris canariensis</i>	1 to 4	9	16
Crabgrass				
Hairy	<i>Digitaria adscendens</i>	2 to 6**	9	16
Large	<i>Digitaria sanguinalis</i>	2 to 6**	9	16
Smooth	<i>Digitaria ischaemum</i>	2 to 6**	9	16
Southern	<i>Digitaria ciliaris</i>	2 to 6**	9	16
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	2 to 6**	9	16
Fall Panicum	<i>Panicum dichotomiflor</i>	2 to 8	9	16
Field Sandbur	<i>Cenchrus incertus</i>	2 to 6	9	16
Foxtail				
Giant	<i>Setaria faberi</i>	2 to 12	9	16
Green	<i>Setaria viridis</i>	2 to 8	9	16
Yellow	<i>Setaria glauca</i>	2 to 8	9	16
Goosegrass	<i>Eleusine indica</i>	2 to 6**	9	16
Itchgrass	<i>Rottboellia cochinchinensis</i>	2 to 6	9	16
Junglerice	<i>Echinochloa colona</i>	2 to 6	9	16
Lovegrass (Stinkgrass)	<i>Eragrostis cilianensis</i>	2 to 6	9	16
Rabbitsfootgrass	<i>Polypogon monspeliensis</i>	1 to 4	9	16
Red Rice	<i>Oryza sativa</i>	1 to 3	9	16
Rygrass				
Hardy	<i>Lolium remotum</i>	2 to 6	9	16
Italian	<i>Lolium multiflorum</i>	2 to 6	9	16
Seedling Johnsongrass	<i>Sorghum halepense</i>	4 to 10	9	16
Shattercane	<i>Sorghum bicolor</i>	6 to 18	9	16
Southwestern Cupgrass	<i>Eriochloa gracilis</i>	2 to 6	9	16
Sprangletop				
Amazon	<i>Leptochloa panicoides</i>	2 to 6	9	16
Bearded	<i>Leptochloa fascicularis</i>	2 to 6	9	16
Mexican	<i>Leptochloa uninervia</i>	2 to 6	9	16
Red	<i>Leptochloa filiformis</i>	2 to 6	9	16

continued

USE DIRECTIONS FOR ANNUAL GRASSES
ALL CROPS (continued)

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT* (inches)	APPLICATION RATES	
			MINIMUM RATE fl oz/A	MAXIMUM RATE ⁽¹⁾ fl oz/A
Texas Panicum	<i>Panicum texanum</i>	2 to 6	9	16
Volunteer Cereals ⁽³⁾				
Barley	<i>Hordeum vulgare</i>	2 to 6	9	16
Oats	<i>Avena sativa</i>	2 to 6	9	16
Rye	<i>Secale cereale</i>	2 to 6	9	16
Wheat ⁽²⁾	<i>Triticum aestivum</i>	2 to 6	9 ⁽²⁾	16
Volunteer Corn ^(2,3)	<i>Zea mays</i>	up to 12	6	12
Volunteer Corn ⁽³⁾	<i>Zea mays</i>	up to 24	9	14
Volunteer Corn ^(2,3)	<i>Zea mays</i>	up to 36	12	16
Volunteer Grain	<i>Sorghum bicolor</i>	8 to 12	9	16
Sorghum				
Wild Oats	<i>Avena fatua</i>	2 to 6	9	16
Wild Proso Millet	<i>Panicum miliaceum</i>	2 to 10	9	16
Witchgrass	<i>Panicum capillare</i>	2 to 8	9	16
Woolly Cupgrass	<i>Eriochloa villosa</i>	2 to 8	9	16

*Generally occurs between 3-leaf stage and tillering.

**Length of lateral growth.

(1) Rates higher than 16 fl oz/A may be applied in certain geographic areas, cropping situations or environmental conditions, where experience has shown that higher rates are needed for satisfactory control of annual grasses. In these situations, rates from 16 to 32 fl oz/A may be applied.

(2) When a cereal grain crop (including wheat) is interseeded for crop establishment or is planted as wind breaks to aid crop establishment, the minimum *Select MAX* Herbicide *with Inside Technology* use rate for control is 12 fl oz/A (0.091 lb ai/A).

(3) Includes Roundup Ready®, Liberty Link® and Clearfield® volunteer corn; however not Sethoxydim-Resistant volunteer corn.

**USE DIRECTIONS FOR PERENNIAL GRASSES
(ALL CROPS)**

- Apply only to actively growing grasses at specified weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the specified growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

Restrictions

- Do not exceed the maximum per application rate listed in CROP SPECIFIC USE DIRECTIONS AND RESTRICTIONS FOR *Select MAX Herbicide with Inside Technology*.
- Do not exceed the maximum yearly rate listed in Table 1, CROP SPECIFIC USE DIRECTIONS AND RESTRICTIONS FOR *Select MAX Herbicide with Inside Technology*.

GRASS SPECIES	WEED HEIGHT (inches)	APPLICATION RATE	
		MINIMUM RATE fl oz/A	MAXIMUM RATE fl oz/A
Bermudagrass (<i>Cynodon dactylon</i>)			
First Application	3 (or up to 6" runners)	12	32
Repeat Application(s) (if regrowth occurs)	3 (or up to 6" runners)	12	32
Fescue, Tall (<i>Festuca arundinacea</i>)			
First Application	4 to 8	12	32
Repeat Application(s) (if regrowth occurs)	4 to 8	12	32
Foxtail Barley (<i>Hordeum jubatum</i>)			
First Application	2 to 6	12	32
Repeat Application (if regrowth occurs)	2 to 6	12	32
Orchardgrass (<i>Dactylis glomerata</i>)			
First Application	4 to 8	12	32
Repeat Application(s) (if regrowth occurs)	4 to 8	12	32
Quackgrass (<i>Elytrigia repens</i>)			
First Application	4 to 12	12	32
Repeat Application(s) (if regrowth occurs)	4 to 12	12	32
Rhizome Johnsongrass (<i>Sorghum halepense</i>)			
First Application	12 to 24	12	32
Repeat Application(s) (if regrowth occurs)	6 to 18	9	24
Wirestem Muhly (<i>Muhlenbergia frondosa</i>)			
First Application	4 to 8	12	32
Repeat Application(s) (if regrowth occurs)	4 to 8	12	32
Perennial Bluegrass [Roughstalk (<i>Poa trivialis</i>)] [Kentucky (<i>Poa prantensis</i>)]			
First Application	2 to 4	12	32
Repeat Application(s)	2 to 4	12	32
Bentgrass (<i>Agrostis</i> spp.)			
First Application	2 to 4	-	32
Repeat Application(s) (if regrowth occurs)	2 to 4	-	32

TANK MIX - LABEL INFORMATION

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture, including all crop rotational and other crop restrictions. Those concerns may include, but are not limited to:

1. Geographic restrictions - all products are not registered for use in all areas and rates may vary from one region of labeled use to another;
2. Crop rotation restrictions;
3. Applicator certification requirements;
4. Worker safety rules (e.g. protective clothing, reentry time, posting);
5. Soil type or soil characteristics (e.g. pH, OM);
6. Maximum dosage or number of applications per year;
7. Rain free period required; or
8. Application timing (e.g. pre-harvest interval)
9. **Restriction:** Do not exceed the total yearly rates.

TANK MIX APPLICATION OF *SELECT MAX* HERBICIDE *WITH INSIDE TECHNOLOGY* AND BROADLEAF HERBICIDES FOR CONTROL OF GRASSES AND BROADLEAF WEEDS

- Apply only to actively growing grass and broadleaf weeds at specified height or growth stage listed on each label.
- Apply when the first grass or broadleaf weed species in a mixed population reaches the specified height or growth stage for treatment.
- Apply under favorable soil moisture and humidity that exist a few days after rainfall or within seven days after irrigation.
- Always add the appropriate adjuvant to the spray mix at the rate prescribed for each specific tank mix combination.
- Tank mix applications may sometimes result in reduced grass control and possible increases in crop injury as compared to either product used alone. If regrowth occurs, or an additional flush of new grass emerges, make a second application of *Select MAX* Herbicide *with Inside Technology*, as specified in the respective size and rate tables.
- **Restriction:** Do not tank mix *Select MAX* Herbicide *with Inside Technology* when broadleaf weeds are tall and/or dense enough to prevent proper grass coverage.

MIXING INSTRUCTIONS

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. While agitating, add the correct amount of *Select MAX* Herbicide *with Inside Technology*. Agitation should create a rippling or rolling action on the water surface.
3. If tank mixing *Select MAX* Herbicide *with Inside Technology* with other labeled herbicides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates and then solutions. Prepare no more spray mixture than is required for the immediate spray operation.
4. Add any required adjuvants (crop oil concentrate, non-ionic surfactant and/or nitrogen solution).
5. Fill spray tank to desired level with water. **Agitation should continue until all spray solution has been applied.**

Failure to agitate the spray solution may result in improper mixing of the herbicides and unsatisfactory weed control. Mixing and compatibility qualities should be verified by a jar test.

PLEASE CONTACT VALENT U.S.A. LLC AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

©2018 Valent U.S.A. LLC

Select MAX and *with Inside Technology* are registered trademark and trademark of Valent U.S.A. LLC

Clearfield is a registered trademark of BASF

Liberty and Liberty Link are registered trademarks of Bayer

Roundup Ready is a registered trademark of Monsanto Company

Manufactured for:
Valent U.S.A. LLC

P.O. Box 8025
Walnut Creek, CA 94596-8025

Made in U.S.A.

2018-SMAX-0010