

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Durango<sup>®</sup> DMA<sup>®</sup>

EPA Reg. No. 62719-556

### Control and Management of Glyphosate-Resistant *Amaranthus* spp. (Not for Use or Distribution Within the State of California)

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

#### Directions for Use

Read the Product Information, Herbicide Resistance Management, Spray Drift Management, Mixing Directions, Application Equipment and Application Methods, and Roundup Ready Crops on the label attached to this product's container.

#### Corn - Preplant, Preemergence, At-Planting

Tank mix this product or apply it sequentially with a herbicide that has a different mode of action to control a naturally occurring glyphosate-resistant biotype of an *Amaranthus* species. To control emerged weeds, apply this product in a tank mix with a preemergence herbicide such as Degree Xtra, Harness Xtra, Harness Xtra 5.6L, or another residual herbicide to control *Amaranthus* spp. Appropriate cultural and mechanical control practices (e.g., crop rotation or tillage) may also be used. For best results, tillage or make a burndown application prior to planting.

#### Corn Hybrids with Roundup Ready<sup>®</sup> 2 Technology – Postemergence (In-Crop)

Tank mix this product or apply it sequentially with a herbicide that has a different mode of action to control a naturally occurring glyphosate-resistant biotype of an *Amaranthus* species. To control emerged weeds, apply this product in a tank mix with a preemergence residual herbicide such as FulTime, Keystone, Keystone LA, SureStart, Surpass EC or TopNotch, or another residual herbicide to control *Amaranthus* spp.

To control emerged weeds as a postemergence application, apply this product in a tank mix with other herbicides such as 2,4-D or dicamba (Clarity, Banvel, or Distinct) and a residual herbicide such as FulTime, Keystone, Keystone LA, SureStart, Surpass EC, or TopNotch for continued control of *Amaranthus* spp. Appropriate cultural and mechanical control practices (e.g., crop rotation or tillage) may also be used. For best results, tillage or make a burndown application prior to planting.

#### Cotton - Preplant, Preemergence, and At-Planting

Tank mix this product or apply it sequentially with a herbicide that has a different mode of action to control a naturally occurring glyphosate-resistant biotype of an *Amaranthus* species. To control emerged weeds, apply this product in a tank mix with a preemergence soil residual herbicide labeled for control of *Amaranthus* spp. such as fluometuron (Cotoran) and/or pendimethalin (Prowl H2O). Appropriate cultural and mechanical control practices (e.g., crop rotation or tillage) may also be used. For best results, tillage or make a burndown application prior to planting.

**Cotton with Roundup Ready® Gene and Roundup Ready® Flex Gene- Postemergence (In-Crop)**

Tank mix this product or apply it sequentially with a herbicide that has a different mode of action to control a naturally occurring glyphosate-resistant biotype of an *Amaranthus* species. To control emerged weeds, apply this product preemergence in a tank mix with metolachlor prior to the 4-leaf stage for the control of *Amaranthus* spp. To control emerged weeds preemergence at layby, apply this product in a tank mix with a residual herbicide such as diuron (Direx) or flumioxazin (Valor) post-directed to control *Amaranthus* spp.

To control emerged weeds as a postemergence application, apply this product in a tank mix with MSMA and diuron (Direx) or flumioxazin (Valor) to control emerged *Amaranthus* spp. Appropriate cultural and mechanical control practices (e.g., crop rotation or tillage) may also be used. For best results, tillage or make a burndown application prior to planting.

**Soybean - Preplant, Preemergence, and At-Planting**

Tank mix this product or apply it sequentially with a herbicide that has a different mode of action to control a naturally occurring glyphosate-resistant biotype of an *Amaranthus* species. To control emerged weeds, apply this product in a tank mix with a preemergence residual herbicide such as alachlor (Intro) or another residual herbicide to control *Amaranthus* spp. For preplant applications, add 2,4-D to the tank mix to help control emerged broadleaf weeds.

To control emerged weeds as a postemergence application, apply this product in a tank mix with lactofen (Cobra) or fomesafen (Flexstar) to control emerged *Amaranthus* spp. Appropriate cultural and mechanical control practices (e.g., crop rotation or tillage) may also be used. For best results, tillage or make a burndown application prior to planting.

®Trademark of Dow AgroSciences LLC

R353-049

EPA accepted 11/26/10

Replaces R353-031