

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Durango[®] DMA[®]

EPA Reg. No. 62719-556

Postemergence Weed Control on Fescue Grown for Seed

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

Directions for Use

Read the General Information, Herbicide Resistance Management, Spray Drift Management, Mixing Directions, and Application Equipment and Application Methods on the label attached to this product's container.

Apply this product on glyphosate-tolerant tall and fine fescue grown for seed only. Use 4 to 12 fl oz of this product per acre as a postemergence spray on glyphosate-tolerant tall fescue selections. Apply six weeks after germination and to established crops after growth resumes in the fall until onset of dormancy, and in the spring after dormancy break until 60 days before harvest.

Applying this product postemergence controls or suppresses the following weeds: annual bluegrass mustards, Canada thistle, cheatgrass, chickweed, dandelion, downy brome, fleabane, pennycress, quackgrass, shepherd's-purse, sowthistle, and wild oat. The specified rate range of this product limits the level of control of certain weed species. **Note:** Some crop discoloration and yellowing may occur at higher rates of application with glyphosate-tolerant tall and fine fescue selections. Reduction in stand of these selections may occur under conditions of stress.

Precautions and Restrictions:

- Do not spray during or within two weeks after periods when the air temperature is below 25°F.
- Remove domestic livestock from the seed production field prior to application.
- Do not resume grazing or harvesting the treated area for 60 days after application.
- Make only two applications of this product per crop growth cycle to any one site. If two applications are required, make only one fall and one spring application during one 12-month cycle.

®Trademark of Dow AgroSciences LLC

R353-036

EPA accepted 11/26/10

Initial printing