

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Durango[®] DMA[®]

EPA Reg. No. 62719-556

Weed Control in Canola (Winter Varieties) with Roundup Ready[®] Gene

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

Directions for Use

Read the General Information, Herbicide Resistance Management, Spray Drift Management, Mixing Directions, Application Equipment and Application Methods, and Roundup Ready Crops on the label attached to this product's container.

Types of Applications: Preplant, at-planting, preemergence, postemergence (in-crop)

Roundup Ready spring canola is defined as Roundup Ready canola varieties that are seeded in the early fall, harvested the following spring or winter, and are intended to enter a cold period dormancy in the winter.

Maximum Allowable Application Rates

Application Type	Rate (pints/acre)
preplant preemergence at-planting	3
total in-crop applications from emergence to canopy closure or prior to bolting in the spring	

Preplant, Preemergent and At-Planting

Apply before, during, or after planting Roundup Ready winter canola.

Precautions and Restrictions:

- Do not apply more than 3 pints of this product per acre per season for all preplant, preemergence and at-planting applications.

Postemergence (In-Crop)

Apply this product as a postemergence application to Roundup Ready winter canola from emergence to canopy closure in the fall and prior to bolting in the spring. Applications made during or after bolting may result in crop injury or yield loss. To maximize yield potential, make applications early to eliminate competing weeds.

A sequential application of this product may be necessary to control some weeds with multiple germination times, suppressed (stunted) weeds, or weeds that have overwintered. Make the second application after some regrowth has occurred and at least 60 days after the previous application of this product.

Single Application: Apply 18 to 24 fl oz per acre in the fall when weeds are small and actively growing. Use the higher rate in the rate range when weed densities are high, when weeds have overwintered, or when weeds become large and well established. Applying more than 18 fl oz per acre prior to the 6-leaf stage may result in reduced crop growth in the fall. Avoid overlapping applications that may result in temporary yellowing and growth reduction.

Sequential Application: Apply 12 to 24 fl oz per acre to 2-leaf or larger canola in the fall followed by a sequential application at the same rate at a minimum interval of 60 days. Make the sequential application before bolting in the spring. Sequential applications are required for early emerging annual weeds and winter emerging weeds such as downy brome, jointed goatgrass and ryegrass, and for weeds that have overwintered. This product controls or suppresses most perennial weeds; however, for some perennial weeds, sequential applications may be required to reduce competition with the crop.

Precautions and Restrictions:

- **Preharvest Interval:** Do not apply within 60 days of harvest.
- Do not apply more than two in-crop (over the top) broadcast applications from crop emergence up to the onset of bolting.
- Do not apply more than 1.5 pints of this product per acre per season for all in-crop applications.
- Reduced crop growth in the fall may result if more than 18 fl oz per acre is applied prior to the 6-leaf stage.
- No waiting period is required between application and open grazing of livestock.

®Trademark of Dow AgroSciences LLC
Roundup Ready® is a registered trademark of Monsanto Company

R353-038
EPA accepted 11/26/10
Initial printing