

Product Bulletin



Chestnut Run Plaza, 974 Centre Road, Wilmington DE 19805, USA

DuPont™ Aproach® Prima Fungicide

EPA Reg. No. 352-883
2(ee) Recommendation†

Control/Suppression of Tar Spot of Corn (*Phyllacora-maydis*)

ATTENTION

† This recommendation is made as permitted under FIFRA 2(ee) and has not been submitted to or approved by the EPA or state lead pesticide agency.

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for DuPont™ Aproach® Prima Fungicide before applying. Carefully follow all precautionary statements and application use directions.
- Use of DuPont™ Aproach® Prima Fungicide according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for DuPont™ Aproach® Prima Fungicide.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

Apply DuPont™ APROACH® PRIMA Fungicide at 3.4-6.8 fl oz. per acre for control of Tar Spot (*Phyllacora maydis*) in corn. Begin applications prior to disease development. Use the highest specified rate and shorter interval when disease pressure is high.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with such non-labeled use.

PRECAUTIONS:

- The minimum re-treatment interval in corn for DuPont™ APROACH® PRIMA Fungicide is 7 days.
- Do not tank mix DuPont™ APROACH® PRIMA Fungicide with an adjuvant or crop oil when spraying corn between V8 and VT stages of growth.
- The minimum pre-harvest interval (PHI) between the last application and harvest for grain or ear is 30-days and for silage is 21-days.
- Do not exceed 6.8 fluid ounces of DuPont™ APROACH® PRIMA Fungicide per acre per year or make more than 2 applications per year.
- When applied alone or in combination with other products containing picoxystrobin, do not apply more than 0.585 pounds of picoxystrobin active ingredient per acre per year to corn.
- When applied alone or in combination with other products containing cyproconazole, do not apply more than 0.036 pounds of cyproconazole active ingredient per acre per year to corn.

® ™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

R601-001

Issued: 04/11/19

Initial printing