

**RESTRICTED USE PESTICIDE**

FOR RETAIL SALE TO AND USE ONLY BY  
CERTIFIED APPLICATORS OR PERSONS UNDER  
THEIR DIRECT SUPERVISION AND ONLY FOR  
THOSE USES COVERED BY THE CERTIFIED  
APPLICATOR'S CERTIFICATION.

**CHLORPYRIFOS 4E AG**

Insecticide

**EPA Reg. No. 66222-19****RESTRICTED USE PESTICIDE  
DUE TO TOXICITY TO FISH AND AQUATIC  
ORGANISMS**

FOR RETAIL SALE TO AND USE ONLY BY  
CERTIFIED APPLICATORS OR PERSONS  
UNDER THEIR DIRECT SUPERVISION AND  
ONLY FOR THOSE USES COVERED BY THE  
CERTIFIED APPLICATOR'S CERTIFICATION.

**SILENCER<sup>®</sup>**

Insecticide

**EPA Reg. No. 66222-104****FIFRA § 2(ee) Recommendation****TANK MIX RECOMMENDATION  
FOR CONTROL OF SOYBEAN APHID IN SOYBEANS.  
(Not registered for use in California or New York)**

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

**DIRECTIONS FOR USE**

It is violation of federal law to use this product in a manner inconsistent with its labeling.

Refer to the Chlorpyrifos 4E AG and Silencer labels for information regarding use restrictions, use precautions, and other information. The most restrictive provisions on either label will apply.

**Foliar Treatment**

Use a tank mix of Chlorpyrifos 4E AG + Silencer at the following rates in sufficient water to ensure thorough coverage of treated plants:

PEST	CHLORPYRIFOS 4E AG	SILENCER <sup>™</sup>
Soybean Aphid	8 ounces/acre	2-3 ounces/acre

Apply as a broadcast spray using either aerial or ground equipment when field counts indicate damaging insect populations are developing or present. When applying by air, apply in a minimum of 2 gals. of water per acre.

Retreat as necessary to maintain control. When heavy populations are present, use the higher rates of each product in the tank mix.

On determinate soybeans, do not apply more than one application after pod set.

**RESTRICTIONS:** Do not apply more than 6 pints (3 lbs. a.i.) of Chlorpyrifos 4E AG per acre per season nor more than 0.48 pint (0.06 lb.a.i.) of Silencer per acre per season. Do not make more than 3 applications per season. Do not apply last treatment within 30 days before harvest nor apply last two treatments closer than 14 days apart. Do not allow livestock to graze in treated areas or otherwise feed treated soybean forage, hay, and straw to meat or dairy animals. On determinate soybeans, do not apply more than one application after pod set.

REV 9-29-09