
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 1070A-S5 0819

Sale, use, and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.

FLUAZIFOP-P-BUTYL	GROUP	1	HERBICIDE
--------------------------	--------------	----------	------------------

Fusilade® DX Herbicide

Postemergence Herbicide for Control of Perennial and Annual Grass Weeds in Agricultural Areas.

This supplemental label expires on 08/21/2022 and must not be used or distributed after this date.

Active Ingredient:

Fluazifop-P-butyl

Butyl (*R*)-2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoate*24.5%

Other Ingredients:75.5%

Total:100.0%

Fusilade® DX Herbicide formulation is an emulsifiable concentrate (EC).

*Fusilade DX Herbicide contains 2 pounds (+) isomer (fluazifop-P-butyl) per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-1070

All applicable directions, restrictions and precautions on the Fusilade DX Herbicide (EPA Reg. No. 100-1070) label are to be followed. Before using Fusilade DX Herbicide as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Use of Fusilade DX Herbicide according to this labeling is subject to the limitations imposed by the label affixed to the container for Fusilade DX Herbicide (EPA Reg. No. 100-1070).

syngenta

DIRECTIONS FOR USE

CROP SPECIFIC RESTRICTIONS

Fusilade DX Rate Conversion Table

fl oz product/acre	lb ai fluazifop/acre
4	0.062
6	0.094
8	0.125
10	0.156
12	0.188
16	0.250
24	0.375
30	0.469
32	0.500
48	0.750
72	1.125

CITRUS FRUITS GROUP 10 (Calamondin; citrus citron; citrus hybrids (includes chironja, tangelo, tangor); grapefruit; kumquat; lemon; lime; mandarin (tangerine); orange, sour; orange, sweet; pummelo; Satsuma mandarin)

- Do not apply a total of more than 72 fl oz of Fusilade DX Herbicide per acre per year (1.125 lb ai/A).
- Do not apply more than 24 fl oz of Fusilade DX Herbicide per acre per application (0.375 lb ai/A).
- Do not apply more than 3 applications per year.
- Maintain a minimum of 21 days between applications.
- Do not harvest within 14 days of last application.

SPECIAL INSTRUCTIONS FOR USE IN APRICOT; BLUEBERRY, LOWBUSH (As well as lingonberry; currant, native); **BUSHBERRY SUBGROUP 13-07B** (Aronia berry; blueberry, highbush; buffalo currant; Chilean guava; cranberry, highbush; currant, black; currant, red; elderberry; European barberry; gooseberry; honeysuckle, edible; huckleberry; jostaberry; Juneberry (Saskatoon berry); salal; sea buckthorn; and cultivars, varieties, and/or hybrids of these); **CANEBERRY SUBGROUP 13-07A** (Blackberry; loganberry; raspberry, black and red; wild raspberry; cultivars, varieties, and /or hybrids of these); **CHERRY; CITRUS FRUITS GROUP 10** (Calamondin; citrus citron; citrus hybrids (includes chironja, tangelo, tangor); grapefruit; kumquat; lemon; lime; mandarin (tangerine); orange, sour; orange, sweet; pummelo; Satsuma mandarin); **MACADAMIA NUT; NECTARINE; PEACH; PECAN; PLUM; PRUNE; SMALL FRUIT VINE CLIMBING**

SUBGROUP 13-07F (EXCEPT FUZZY KIWIFRUIT) (Amur River grape; gooseberry; grape; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and/or hybrids of these)

Apply Fusilade DX Herbicide as a directed postemergence application in the interspaces and around the base of plants while avoiding contact of spray with plants. Apply with sufficient spray volume and pressure to ensure complete coverage of target grasses. Applications can be made as a broadcast, strip band, or spot spray application at rates and growth stages listed in Tables 2-5. Refer to the Spray Additives section for spray additives.

Fusilade® and the Syngenta logo are trademarks of a Syngenta Group Company.
©2019 Syngenta