
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 1313A-S10 0218

AZOXYSTROBIN	GROUP	11	FUNGICIDE
DIFENOCONAZOLE	GROUP	3	FUNGICIDE

Quadris Top® Fungicide

This supplemental label expires on 02/16/2021 and must not be used or distributed after this date.

Active Ingredients:

Azoxystrobin*	18.2%
Difenoconazole**	11.4%
Other Ingredients:	70.4%
Total:	100.0%

*CAS No. 131860-33-8

**CAS No. 119446-68-3

Quadris Top Fungicide is formulated as a suspension concentrate (SC) containing 1.67 lb of azoxystrobin active ingredient and 1.05 lb of difenoconazole active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use on container label.

EPA Reg. No. 100-1313

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Quadris Top Fungicide as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
Cranberry	Bitter rot <i>(Colletotrichum gloeosporioides)</i> Blotch rot <i>(Physalospora vacciniae)</i> Cottonball <i>(Monilinia oxycocci)</i> Fruit Rots <i>(Physalospora vacciniae)</i> <i>(Glomerella cingulata)</i> <i>(Coleophoma empetri)</i> Leaf rust <i>(Pucciniastrum vaccinii)</i> Lophodermium Twig Blight <i>(Lophodermium spp.)</i> Ripe rot <i>(Coleophoma empetri)</i>	10 – 14*	For best activity, apply Quadris Top Fungicide prior to or early in the disease development. An adjuvant may be added at specified rates. Apply on a 7-14 day interval. Do not apply more than two sequential applications before alternating to a fungicide with a different mode of action.

*10-14 fl oz product/A contains 0.082-0.115 lb difenoconazole/A and 0.131-0.183 lb azoxystrobin/A.

Application: For best results, sufficient water volume must be used to provide thorough coverage. Quadris Top Fungicide can be applied by ground, chemigation, or aerial application. For aerial applications, apply in a minimum of 5 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy. Applicators should use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) Do not apply more than 42 fl oz/A/year of Quadris Top Fungicide.
- 2) Do not apply more than 0.34 lb ai/A/year of difenoconazole-containing products.
- 3) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 4) Do not allow release of irrigation or flood water to non-target aquatic habitat for at least 14 days after the last application.
- 5) Do not apply when weather conditions favor drift from treated areas to non-target aquatic habitat.
- 6) Do not treat fields used for aquaculture of fish or crustacean.
- 7) Do not drain water from treated fields into ponds used for aquaculture of fish or crustacean.
- 8) Do not use water drained from treated field to irrigate other crops.
- 9) Do not apply to flooded crop.
- 10) Do not apply Quadris Top Fungicide within 30 days of harvest (30-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
Guava	<p>Alternaria Fruit Rot</p> <p>Anthracnose (<i>Colletotrichum gloeosporioides</i>)</p> <p>Suppression Rust (<i>Puccinia psidii</i>)</p>	10 – 14*	<p>For best activity, apply Quadris Top Fungicide prior to or early in the disease development. An adjuvant may be added at specified rates. Apply on 10-14 day interval.</p> <p>Do not apply more than two sequential applications before alternating to a fungicide with a different mode of action.</p>

*10-14 fl oz product/A contains 0.082-0.115 lb difenoconazole/A and 0.131-0.183 lb azoxystrobin/A.

Application: For best results, sufficient water volume must be used to provide thorough coverage. Quadris Top Fungicide can be applied by ground, chemigation, or aerial application. For aerial applications, apply in a minimum of 10 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy. Applicators should use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) Do not apply more than 56 fl oz/A/year of Quadris Top Fungicide.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 4) Quadris Top Fungicide may be applied the day of harvest (0-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
Papaya	<p>Blossom blight and fruit rot (<i>Colletotrichum gloeosporioides</i>)</p> <p>Alternaria fruit spot (<i>A. alternata</i>)</p> <p>Powdery Mildew (<i>Oidium</i> spp.)</p> <p>Brown Spot (<i>Corynespora cassicola</i>)</p>	10 – 14*	<p>For best activity, apply Quadris Top Fungicide prior to or early in the disease development. An adjuvant may be added at specified rates. Apply on 10-14 day interval.</p> <p>Do not apply more than two sequential applications before alternating to a fungicide with a different mode of action.</p>

*10-14 fl oz product/A contains 0.082-0.115 lb difenoconazole/A and 0.131-0.183 lb azoxystrobin/A.

Application: For best results, sufficient water volume must be used to provide thorough coverage. Quadris Top Fungicide can be applied by ground, chemigation, or aerial application. For aerial applications, apply in a minimum of 10 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy. Applicators should use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) Do not apply more than 56 fl oz/A/year of Quadris Top Fungicide.
- 2) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 4) Quadris Top Fungicide may be applied the day of harvest (0-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
Watercress	Cercospora leafspot (<i>Cercospora</i> spp.)	10 - 14	For best activity, apply Quadris Top Fungicide prior to or early in the disease development. An adjuvant may be added at specified rates. Apply on a 7-14 day interval. For applications made to watercress, production fields must be drained of water at least 24 hours prior to application and water must not be reapplied to the field for a minimum of 24 hours following the application. Do not apply more than two sequential applications before alternating to a fungicide with a different mode of action.

Application: For best results, sufficient water volume must be used to provide thorough coverage. Quadris Top Fungicide can be applied by ground, chemigation, or aerial application. For aerial applications a minimum of 5 gal/A of water is recommended. Applicators should use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) Do not apply directly to water and do not allow water in a treated field for at least 24 hours.
- 2) Do not apply more than 56 fl oz/A/year of Quadris Top Fungicide.
- 3) Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 4) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 5) Do not apply more than 0.75 lb ai of azoxystrobin-containing products per acre per cutting.
- 6) Do not apply Quadris Top Fungicide within 30 days of harvest (30-day PHI).

©2018 Syngenta

Quadris Top® and the Syngenta logo are trademarks of a Syngenta Group Company