



# Sevin® XLR PLUS

brand Carbaryl Insecticide

EPA Reg. No. 61842-37

For Control of Spotted Lanternfly  
Scientific Name: (*Lycorma delicatula*)

For use on Grapes, Pome Fruit and Stone Fruit

## 2(EЕ) RECOMMENDATION - USE INFORMATION IN ACCORDANCE WITH SECTION 2(EЕ) OF FIFRA

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This recommendation must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

CROP	Pest	RATE OF APPLICATION – Quarts Per Acre	SPECIFIC DIRECTIONS
Grapes	Spotted Lanternfly ( <i>Lycorma delicatula</i> )	2	OBSERVE BEE CAUTION*** Repeat applications as necessary up to a total of 5 times per year but not more often than once every 7 days. Do not concentrate spray on the bunch or visible residues may result.
Pome Fruit		3	OBSERVE BEE CAUTION*** On apples, avoid use during the period from full bloom until 30 days after full bloom unless fruit thinning is desired. Use for pesticial activity during this period may also result in fruit removal. Repeat applications as necessary up to a total of 8 times per crop (including thinning sprays on apples) but not more often than once every 14 days.
Stone Fruit		3	OBSERVE BEE CAUTION*** Repeat applications as necessary up to a total of 3 times per crop but no more often than once every 7 days. An additional application at the dormant or delayed dormant timing may be made.

\*\*\* **BEE CAUTION:** This product is highly toxic to honeybees and other bees exposed to direct treatment or residues on crops or weeds in bloom. This product may show residual toxicity to honeybees, especially in humid climates and under slow drying conditions. Notifying beekeepers within 1 mile of treatment area at least 48 hours before product is applied will allow them to take additional steps to protect their bees. Limiting application to times when bees are least active, e.g., within 2 hours of sunrise or sunset, will minimize risk to bees. Do not apply this product to target crops or weeds in bloom. If weed or cover crop bloom is present mow orchard floor or between rows prior to applying this product.

### RESTRICTIONS: GRAPES

**East of the Rocky Mountains**, the restricted-entry interval (REI) = 2 days. **EXCEPTION:** the REI is 6 days for grape girdling and cane turning.

**West of the Rocky Mountains**, the restricted-entry interval (REI) = 6 days.

TESSENDERLO KERLEY, INC., member of TESSENDERLO GROUP

P. O. Box 15627, Phoenix, Arizona 85060-5627  
2255 North 44<sup>th</sup> Street, Suite 300, Phoenix, Arizona 85008-3279  
Tel. (602) 889-8300, Fax (602) 889-8430



**RESTRICTIONS: POME FRUIT**

- Do not apply to quince.
- Do not use on pears between the tight flower cluster up to the 20 mm fruit size. Use during this period may result in undesirable fruit thinning and/or deformed fruit.
- Do not apply within 3 days of harvest.
- Do not apply more than a total of 15 quarts per acre per crop per year.
- Do not make more than a total of 8 applications per crop per year.

**RESTRICTIONS: STONE FRUIT**

- Do not apply within 3 days of harvest, except in California. In California, do not apply within 1 day of harvest.
- All States including California:  
Do not apply more than a total of 14 quarts per acre per crop per year.  
Do not apply more than a total of 5 quarts per acre at the dormant or delayed dormant timing.  
Do not apply more than a total of 9 quarts per acre during the production season.
- California only: Do not apply more than 4 quarts per acre per application during the production season.
- All States other than California: Do not apply more than 3 quarts per acre per application during the production season.

**RESTRICTIONS AND PRECAUTIONS**

- Carefully read and follow all mixing and handling instructions on the EPA registered label.
- Do not allow public use of treated areas during application or until sprays have dried.
- Avoid direct application to lakes, streams and ponds.
- Do not directly spray or allow spray drift to contact surfaces where visible spray deposit may be objectionable.

**IMPORTANT NOTICE**

This bulletin contains new or supplemental instructions for the use of this product which do not appear on the EPA-registered package label. Follow instructions carefully. This labeling must be in the possession of the user at the time of pesticide application. Read the Limitation of Warranty and Liability on the product label before buying or using this product. If terms are not acceptable, return the unopened packaging at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the product label.

For MEDICAL Emergencies Call 24 Hours A Day 1-866-374-1975. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

Tessenderlo Kerley, Inc.  
2255 N. 44th Street, Suite 300  
Phoenix, AZ 85008 USA  
1-800-525-2803  
[www.novasource.com](http://www.novasource.com)

NovaSource and Sevin are registered trademarks of Tessenderlo Kerley, Inc. ©2012, 2018 Tessenderlo Kerley, Inc. All rights reserved.  
SEVIN-XLRPLUS-rev112018