RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.





ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

This recommendation for CAPTURE LFR Insecticide is valid until 03/18/2025, or until withdrawn, cancelled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

CROP	PEST	RATE OF APPLICATION
Corn (At-Plant) including: Field corn, Popcorn, Field corn grown for seed, Sweet corn, Sweet corn grown for seed	Asiatic Garden Beetle	8.5 fl oz/A (0.1 lb ai/A)

DIRECTIONS FOR USE

- To reduce corn plant stand loss, apply Capture LFR Insecticide at-plant as a 5 to 7 inch band (T- band) over the open furrow, or in-furrow with the seed.
- May be tank mixed with other pesticides and fertilizers.

RESTRICTIONS:

- Do not apply more than 0.1 lb ai/A (8.5 fl oz/A) per season as an at-plant application.
- For Field Corn Do not apply more than 0.3 lb ai/A per season including pre-plant incorporated, atplant, pre-emergence, and foliar applications of other bifenthrin products.
- For Sweet Corn Do not apply more than 0.2 lb ai/A per season including pre-plant incorporated, atplant, pre-emergence, and foliar applications of other bifenthrin products (such as BRIGADE 2EC Insecticide/Miticide).

