

**RESTRICTED USE PESTICIDE
DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS.
For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and
only for those uses covered by the Certified Applicator's certification.**



FIFRA Section 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF NEW YORK



EPA Reg. No. 59639-35

TO CONTROL AN ADDITIONAL PEST, SPOTTED LANTERNFLY

This FIFRA section 2(ee) recommendation expires on December 31, 2023.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This 2(ee) recommendation contains directions for use which do not appear on the package label and must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions and precautions on the EPA registered label when using *Danitol* 2.4 EC Spray. Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

| TO CONTROL AN ADDITIONAL PEST, SPOTTED LANTERNFLY | | |
|---|--|---|
| Crop | Application Rates | Special Instructions |
| <p style="text-align: center;">Grape</p> <p>Do not apply within 21 days of harvest. Do not exceed 2-2/3 pts (42-2/3 fl oz, 0.8 lb ai) total application of <i>Danitol</i> 2.4 EC Spray per acre per season.</p> | <p>16 to 21-1/3 fl oz/A (0.3 to 0.4 lb ai/A)</p> | <p>Make foliar applications of <i>Danitol</i> 2.4 EC when adults are actively flying in the perimeter and within the orchard.</p> <p>Apply by ground with airblast equipment as a full coverage spray or apply by air in a minimum of 10 gals/A and repeat as needed to maintain control but not more often than every 10 days.</p> |
| <p style="text-align: center;">Pome Fruit</p> <p>Do not apply within 14 days of harvest. Do not exceed 2-2/3 pts (42-2/3 fl oz, 0.8 lb ai) total application of <i>Danitol</i> 2.4 EC Spray per acre per season. Do not apply as a ULV spray. Do not feed or allow livestock to graze on cover crops from treated orchards.</p> | | <p>Resistance Management: It is generally recommended that no more than two <i>Danitol</i> 2.4 EC Spray applications be made per season as part of a Resistance Management Program. Use non-pyrethroid products at other timings to control pests.</p> |
| <p style="text-align: center;">Stone Fruit</p> <p>Do not apply within 14 days of harvest. Do not exceed 2-2/3 pts (42-2/3 fl oz, 0.8 lb ai) total application of <i>Danitol</i> 2.4 EC Spray per acre per season. Do not apply as a ULV spray. Do not feed or allow livestock to graze on cover crops from treated orchards.</p> | | |

Danitol is a registered trademark of Sumitomo Chemical Company, Ltd.

Registrant: Valent U.S.A. LLC
P.O. Box 8025
Walnut Creek, CA 94596-8025