

RESTRICTED USE PESTICIDE
DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS.
 For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

FIFRA Section 2(ee) Recommendation



THIS RECOMMENDATION IS ONLY FOR THE STATES OF MAINE,
 MASSACHUSETTS, MICHIGAN, NEW HAMPSHIRE, NEW JERSEY,
 OREGON, PENNSYLVANIA, VIRGINIA, WASHINGTON, WEST VIRGINIA



EPA Reg. No. 59639-35

DANITOL® 2.4 EC SPRAY FOR USE ON BLACK STEM BORER IN POME FRUIT

This FIFRA section 2(ee) recommendation expires on December 31, 2024.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This 2(ee) recommendation contains directions for use which do not appear on the package label and must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions and precautions on the EPA registered label when using *Danitol* 2.4 EC Spray. Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

POME FRUIT		
(Including Apple, Crabapple, Loquat, Mayhaw, Quince, Quince (Chinese), Quince (Japanese), Tejocote, Cultivars, varieties, and/or hybrids of these.		
Pest	<i>Danitol</i> 2.4 EC Spray Application Rates	Special Instructions
Black Stem Borer	16 to 21-1/3 fl oz/A (0.3 to 0.4 lb ai/A)	<p>Make foliar applications of <i>Danitol</i> 2.4 EC when black stem borer adults are actively flying in the perimeter and within the orchard.</p> <p>Apply by ground with airblast equipment as a full coverage spray or apply by air in a minimum of 10 gallons per acre and repeat as needed to maintain control but not more often than every 10 days.</p> <p>Resistance Management: It is generally recommended that no more than two <i>Danitol</i> 2.4 EC Spray applications be made per season as part of a Resistance Management Program. Use non-pyrethroid products at other timings to control pests.</p>

Danitol is a registered trademark of Sumitomo Chemical Company, Ltd.

Registrant: Valent U.S.A. LLC
 P.O. Box 8025
 Walnut Creek, CA 94596-8025