RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.



FIFRA 2(ee) Recommendation EPA Reg. No. 279-3426

FOR DISTRIBUTION AND USE ONLY IN DE, GA, IL, ME, MD, MA, MI, NJ, NC, OH, OR, PA, SC, VA, WV AND WI

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

This recommendation for MUSTANG MAXX Insecticide is valid until 03/26/2025, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

| CROP | PEST | RATE OF APPLICATION |
|--|--|--------------------------------|
| Berries Crop Group including: Blackberry Loganberry Red and Black Raspberry Highbush and Lowbush Blueberry Currant | Cherry Fruitworm Cranberry Fruitworm Cranberry Weevil, Japanese Beetle, Plum Curculio Scale (crawlers) | 4.0 fl oz/A (0.025 lb ai/A) |
| Elderberry | Suppression Only: | |
| Gooseberry | Green Aphid, | |
| Huckleberry | Blueberry Aphid, | |
| And cultivars and/or hybrids of these | Blueberry Maggot, | |

DIRECTIONS FOR USE:

Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Thorough coverage is essential to achieve control. Do not exceed maximum allowable rate.

Apply by air or ground equipment using sufficient water to obtain full coverage of foliage. When applying by ground, apply in a minimum of 20 gallons of water per acre. When applying by air, apply in a minimum of 2 gallons of water per acre.

RESTRICTIONS:

- Do not apply more than 4 fl oz/A (0.025 lb ai/A) per application.
- Do not apply more than 24 fl oz/A (0.15 lb ai/A) per season.
- Do not make application less than 7 days apart.
- PHI: 1 Day

